

Dairy cows under the influence of dietary manipulations: A-Review**Asad Ali Khaskheli^{1*}**¹Department of Animal Nutrition, Sindh Agriculture University, Tando jam, Pakistan***Corresponding Author:** Asad Ali Khaskheli, Department of Animal Nutrition, Sindh Agriculture University, Tando jam, Pakistan, E-mail: khaskheliias@gmail.com**Citation:** Dairy cows under the influence of dietary manipulations: A-Review, Sci J of Ani and Vet Sci.2020; 2(1): 01-07.**Submitted:** 13 July 2020; **Approved:** 11 July 2020; **Published:** 14 July 2020**Abstract:**

This review study was carried out in order to understand the dietary influence on the dairy cows. The results obtained from review of already conducted studies revealed that dairy cow is significantly influenced by composition, quality, amount and regime of diet. Maximum daily milk production, milk protein, milk lactose, milk fat, total solids are recorded in dairy when ad-libitum feed and water is provided. Further, water is necessary for maintaining body fluids and proper ion balance; digestion, absorption, metabolization of nutrients; elimination and body cooling. Feeding and water frequency stimulates mammary functions and milk synthesis is manipulated specifically by a non-invasive method. Reducing feeding frequency from 2x daily to 1x daily decreased milk yield from 7 to 38% in dairy cows, however changing feeding frequency from 2x to 3x resulted 18% increase in milk production. In conclusion, diet play a key role in the life of dairy. Good quality feed and water with on ad libitum supports performance and production of dairy cow.

Keywords: Ad Libitum, Diet, Performance, Production**Introduction**

Dairy cow responses to various types of diets differently. Dairy farmers can use knowledge of animal behavior to improve cow well-being and yield. For instance, feeding and watering systems must be placed appropriately. Accessibility of feed and water may be more important than the actual amount of nutrients provided.¹ Efforts must be made to reduce the competition for feed, water, minerals, and shelter. Also, cow space, cow density, and distribution of feed and water are closely related factors. Feed intake and consequent milk yield are improved by provision of feed on cows need and want to eat.² When one cow eats, another might be stimulated to do likewise, whether she is hungry or not. This behavior is an example of social facilitation when cows eat in groups, they eat more than when they are fed separately. Furthermore, cows kept in groups are likely to be less fearful, and hence, more contented,

healthier, and more productive. The common practice of feeding and milking cows in groups thus has a sound psychological basis.³

Water is an essential component to sustain life and optimize growth, lactation, and reproduction of dairy cattle. However, unlike the careful and continuous attention paid by dairy producers and nutritionists to other nutrients in the ration, oftentimes the quality and provision of free drinking water does not receive the attention necessary to ensure optimal nutrition and cattle performance. The water requirement per unit of body mass of a high-producing dairy cow is greater than that of any other land-based mammal.⁴ This is because of the high yield of a secretion that is 87% water. Water also is required for digestion and metabolism of energy and nutrients; transport in circulation of nutrients and metabolites to and from tissues; excretion of waste products (via urine, feces, and respiration); maintenance of proper ion, fluid,

Cite this article: Asad Ali Khaskheli, Dairy cows under the influence of dietary manipulations: A-Review, Sci J of Ani and Vet Sci.2020; 2(1): 01-07.

and heat balance; and, as a fluid and cushioning environment for the developing fetus.⁵ Total body water content of the bodies of adult dairy cattle ranges between 56 and 81% of body weight depending upon stage in the lactation cycle. Loss of only about 20% of total body water is fatal. Further, it was reported that milk production increases with increment of water intake.⁶ An other scientist reported that the reducing or decreasing watering had negative effect on milk production whereas free access to drinking water had a positive effect on milk yield.⁷

Dairy cows must consume required quantities of water and feed for the production of milk. The amount of water that a cow will drink depends largely upon the ambient temperature, the kind of feed consumed, milk yield and the temperature and cleanliness of the water. Voluntary water intake of cows for optimum milk production depends upon frequent access, without discomfort and at a moderate temperature.⁸ Most cattle normally consume 3 to 4 units of water for each unit of dry feed. Water requirements are directly related to dry matter intake i.e. increased consumption of dry matter will increase water consumption and vice versa. Milking cows require 4-5 kg of water for each kg of milk produced.⁹ Keeping in view the importance of diet in life of dairy cow current study was planned, whereby the main objective was to understanding dietary influence on dairy cow worldwide.

Influence of dietary water on dairy cows

Water constitutes 60 to 70 percent of the body of dairy cow. Water is necessary for maintaining body fluids and proper ion balance; for digesting, absorbing, and metabolizing nutrients; for eliminating waste material and excess heat from the body; for providing a fluid environment for the fetus; and for transporting nutrients to and from body tissues. The water that dairy cattle need is supplied by drinking, by the feed that they consume, and by metabolic water produced by the oxidation of organic nutrients. Water loss occurs via saliva, urine, feces, and milk; through sweating; and by evaporation from body surfaces and the respiratory tract. The amount of water lost from the body of cattle is influenced by the activity of the animal, air temperature, humidity, respiratory rate,

water intake, feed consumption, milk production and other factors.¹⁰

The water requirement per unit of body mass of the high-producing dairy cow is higher than that of any other land-based mammal. This is because she produces a large amount of milk which is 87% water.¹¹ Water is required for digestion and metabolism of energy and nutrients, transport of nutrients and metabolites to and from cells in blood, excretion of waste products (via urine, feces, and respiration), maintenance of proper ion, fluid, and heat balance, and, as a fluid environment for the developing fetus.¹² Total body water content of adult dairy cattle ranges between 56 and 81% of body weight.¹³ Cows in early lactation have more live body weight as water compared with cows in later lactation (69.0 vs. 62.4%); body water content of late pregnant dry cows was 65% of total body weight.¹⁴ About two-thirds of water in the cow's body is in the intracellular compartment. The remaining one-third of water is in extracellular spaces around cells and connective tissues, in blood, and as transcellular water or water in the digestive tract. Water in the digestive tract accounts for 15 to 35% of body weight.¹⁵ About 15% of body weight was as water in the digestive tract of dairy cows in early lactation; in late lactation and during the dry period about 10% of body weight was water in the digestive tract.¹⁶

Another researcher determined that the residence time of a molecule of water in the rumen of lactating dairy cows was about 1 hour. Loss of about 20% of total body water is fatal. Loss of water from the body occurs through milk production, urinary and fecal excretion, sweating, and evaporative loss from the lungs.¹⁷ Daily water losses via milk secretion (73 lb/cow per day) represented between 26 and 34% of total water intake (drinking water plus water in feed consumed).¹⁸ Water lost in feces of lactating cows ranged from 30 to 35% of total water intake, whereas losses in urine were 15 to 22%. Fecal water losses are increased by increasing dry matter intake (DMI), dry matter (DM) content of the diet, and with increasing forage content of the diet. Urinary water excretion is related positively with water availability, amount of water absorbed from the digestive tract, urinary nitrogen, sodium, and potassium

Cite this article: Asad Ali Khaskheli, Dairy cows under the influence of dietary manipulations: A-Review, Sci J of Ani and Vet Sci.2020; 2(1): 01-07.

excretion, and negatively related with dietary DM content.¹⁹ Further it was calculated that losses associated with sweat, saliva, and respiratory evaporation accounted for about 18% of total water loss within the thermoneutral zone. However, amounts and proportions of water loss associated with these routes were highly dependent upon environmental temperature.²⁰

Several factors influences the daily water requirements and intake by dairy cows, including physiological state, amount of milk yield, amount of feed intake, body size, level and kind of activity, environmental factors such as temperature and air movement, diet composition including types of feedstuffs (e.g., concentrate, fresh forages, fermented forages, and hays) as well as nutrient composition (e.g., dietary sodium, potassium, and crude protein contents), and quality (or anti-quality) factors in a particular water source. Other factors affecting consumption may include frequency and periodicity of watering, temperature of the water, and social and behavioral interactions of animals. Water requirements of dairy cattle are met mainly from that ingested as drinking (free) water; that found in or on feed consumed, and, a small amount from metabolic oxidation (metabolic water). For all practical purposes drinking water intake plus that associated with the ration represent total water consumption. Seventy to 97% of total water consumption by lactating dairy cows was from drinking water.²¹ Dry matter content of the diet also is an important factor affecting total water consumption. In totally mixed rations with DM contents ranging from 50 to 70%. Further it was found relatively small differences in drinking water intake; however, when dietary DM content declined from 50 to 30% (ration moisture content increased from 50 to 70%), drinking water intake declined by 42%.²² Moreover, estimating drinking water intake of lactating dairy cows on pasture was studied and it was noticed that that only 38% of total water consumption is free drinking water. Diets with high amounts of sodium-containing salts (e.g., NaCl, NaHCO₃) or protein stimulate water intake.²³

In other study, it has been stated high dry forage diets also may increase water requirements because of higher excretion

of water in feces compared with lower forage diets. There is a direct relationship between DMI and water intake in cattle. If water intake is sub-normal, feed DM intake typically will decrease. However, if water intake is normal and sufficient to meet the physiological needs of the animal for maintenance, growth, lactation and pregnancy, there is no evidence to suggest that increasing water intake (e.g., forced-hydration) beyond normal will result in greater feed DMI or performance.²⁴ Factors typically considered in water quality evaluation include odor and taste, physical and chemical properties, presence of toxic compounds, concentration of macro- and micro-minerals, and microbial contamination. These factors may have direct effects on the acceptability (palatability) of drinking water, or they may affect the animal's digestive and physiological functions, once consumed and absorbed.²⁵ Primary anti-quality factors known to affect dairy cattle include total dissolved solids, sulfur, sulfate and chloride (both being anions), nitrates, iron, and fluoride. Many other potential factors typically listed in water analyses reports and listed as potential risks for humans have not been well-documented in the research literature or under practical conditions to affect dairy cattle performance or health; examples include pH of water (pH between 6 and 9 is assumed acceptable and has very little influence on ruminal pH due to the highly reductive environment in the rumen), total hardness, calcium and magnesium contents.²⁶ It is always possible that isolated cases of higher than normal concentrations of mineral elements, microorganisms, or other toxic compounds may be present and deleterious to cattle. However, typically these cases are extremely difficult to identify and to prove cause and effect.²⁷

Influence of feeding on dairy cows

Regular and complete milking is one of the requirements for continuance of lactation. The effect of changes in feeding frequency on milk yield varies widely between individual species. Reducing feeding frequency from 2x daily to 1x daily decreased milk yield from 7 to 38% in dairy cows, 15 to 48% in ewes, 6 to 35% in dairy cows and increased the rate of loss of udder tissue.²⁸ However increasing milking frequency from 2x to 3x increased milk yield by 7 to 20%.²⁹ The mechanisms responsible

for the increase in milk yield have not been identified but some researchers suggest an increase in mammary epithelial cell (MEC) number, reduction in MEC apoptosis, increased cell activity and frequent removal of feedback inhibitor of lactation (FIL) from the glands.³⁰

Feeding frequency stimulates mammary functions and milk synthesis is manipulated specifically by a non-invasive method. These increases are achieved with little loss of body weight or condition, the extra nutrient requirement being met by an increased feed intake. Significant increase in milk yield with increased feeding frequency.³¹⁻³² The mechanisms responsible for the increase in milk yield include, increase in mammary epithelial cell (MEC) number, reduction in MEC apoptosis, increased cell activity and frequent removal of FIL from the glands. Milk yield for 1x daily milking was significantly lowered ($p<0.05$) observed with 2x and 3x daily milking.³³⁻³⁴ While another scientist reported that changing feeding frequency from 2x to 3x in cattle resulted in 18% increase.³⁵ Some other reported an increase of 10- 20% in cows' milk when daily feeding frequency was changed from 2x to 3x. Further it has been studied that diet during lactation had significant effect ($p<0.001$) on milk yield. Yield declined by 2.5 mL for each additional day of lactation.³⁶⁻³⁷ This result agrees with that of another scientist who reported that milk production in cows gradually declines after reaching its peak.³⁸ This decline was mainly due to the loss of secretory tissue and decrease in rate of secretion per cell.³⁹ Similarly, in non pregnant cows, the decline in milk yield after peak lactation was very gradual, with each month's yield being 95% of preceding month. The study revealed that feeding frequency had significant on milk yield. Yield declined by 2.5 mL for each additional day of lactation. This result agrees with that of who reported that milk production in cows gradually declines after reaching its peak.⁴⁰ This decline was mainly due to the loss of secretory tissue and decrease in rate of secretion per cell. Similarly, in non pregnant cows, the decline in milk yield after peak lactation was very gradual, with each month's yield being 95% of preceding month. It was also reported that cows milked 6x daily consumed higher dry matter than did cows milked 3x daily.⁴¹

In another study a researcher stated that a measurable increase in dry matter intake in cows accompanied the rise in milk production from higher milking frequencies. It was noticeable that as the day of lactation progressed, most of the animals did not gain weight. In fact, some actually lost weight during the milking period (especially animals on 3x daily milking) indicating higher demands for dry matter intake at higher milking frequency.⁴² This result corroborates the findings of another researcher who reported that increased dry matter intake by cows milked 6x daily did not compensate for the increased energy demands, thus these cows lost BW, had a lower body condition scores during the initial lactation period and displayed a longer recovery period than did cows milked 3x daily.⁴³ It has also been reported that cows milked 3x daily did tend ($p<0.1$) to be lighter than their counterparts milked 2x daily during lactation. They observed further, that dry matter intake increased by approximately 15% and cows on 2x and 3x daily milking consumed 10 and 11% more dry matter than controls during lactation respectively.⁴⁴ Although, a researcher reported that increased dry matter intake was too small to be measurable or nonexistent. Further it was reported that cows milked 3x daily actually had lower dry matter intake than those milked 2x daily.⁴⁵

In another study a researcher has reported that milking 3x daily reduced body weight gains in cows and there was a tendency for goats milked 3x daily to lose weight in the present study. Even though, cows were allowed ad libitum dry matter intake, the cows milked 3x daily did not respond to the increased energy demand associated with increased yield by consuming more ration. It appears that increasing feeding frequency results in either preferential utilization of nutrients for milk production or a higher rate of tissue catabolism.⁴⁶ Another researcher suggested that increased milk yield and milk yield: feed intake occurs only as long as tissue reserves are able to subsidize the nutrient needed for milk yield. Day of lactation had significant effect ($p<0.001$) on dry matter intake increasing by 3.73 g for each additional day of lactation.⁴⁷ These results agree with those of scientist who reported that the onset of lactation results in a dramatic increase

in the requirements of nutrients, for example glucose, amino acids and fatty acids. This increase in requirements is met partly by increased voluntary intake partly by an array of metabolic adaptations.⁴⁸ According to another scientist, changes which are of major importance for the establishment and maintenance of high milk production include hypertrophy of the gastro intestinal tract, increased fatty acid metabolism from adipose tissue and an increased rate of gluconeogenesis. The efficient utilization of feed by the animals could be attributed to the breed, physiological status of the animals and the quality of the feed offered.⁴⁹ This observation agrees with the findings by other researchers who reported that feed efficiency for milk production depends on diet, environmental factors and on genetic ability of the animal to utilize these inputs to produce milk.⁵⁰ Moreover, few others noted that efficiency of carbohydrate utilization (carbohydrates being the major nutritive portion of ruminant animal feed) can be increased by treatments which encourage the animals to produce propionate rather than acetate or butyrate from carbohydrates. Consequently, if the animal is making more propionate, it will be found to be using its feed more efficiently. The elephant grass offered to the animals was chopped and fresh thereby increasing the surface area of the roughage.⁵¹ This may therefore, contribute to its efficient utilization by the animals. According to some other scientists, the hormones (e.g. growth hormone, insulin, prolactin) interact to control partitioning of dietary energy into milk and body tissue; this interaction is associated with genetic differences for milk within and between breeds. Further it has been stated the animals could also be under favourable hormonal interaction.⁵²⁻⁵³ The result also revealed that the higher the milking frequency, the higher the efficiency of feed utilization for milk production [1x (0.237), 2x (0.412) and 3x (0.431)]. Animals on 3x milking frequency utilized their feed better than 2x and 1x milking frequencies. This result agrees with those of Barnes et al. (1990) who reported that cows milked 3x daily had efficiency ratios approximately 14% greater than for cows milked 2x. In the present study, animals on 3x daily milking frequency had efficiency ratio of 4.6% greater than 2x milking frequency.

Therefore, animals on 3x milking frequency could have performed better if they were fed according to yield. This could probably explain the non significant difference recorded between 3x and 2x milking frequency in this study.⁵⁴

Conclusions and suggestions

Present study concludes that the diet including water and feed play the key role for normal physiological processes, performance and production of dairy cow. It is further concluded that dairy performance significantly higher high quality diet is provided on ad-libitum. Further studies should be conducted on the dietary effects on the blood biochemistry, immune response and molecular aspects of dairy cows.

References

1. Wilde C.J., Henderson A.J., Knight C.H., Blatchford D.R., Faulkner A. and Vernon R.G., Effects of long-term 3x daily milking on mammary enzyme activity, cell population and milk yield in the goat. *Journal of Animal Science.*, 64: 533-539 (2009).
2. VanBaale M.J., Ledwith D.R., Thomson J.M., Burgos R., Collier R.J. and Baumgard L.H. Effect of increased milking frequency in early lactation with or without recombinant bovine somatotropin. *Journal of Dairy Science.*, 88: 3905-3912 (2015).
3. Thokal M.R., Patil V.C. and Udar S.A., Effect of drinking water frequency on milk yield, fat, total solids and solids-not-fat content in crossbred cows. *Indian Journal of Animal Research.*, 38: 47-49 (2004).
4. Tyrrell H.F., Brown A.C.G., Reynolds P.J., Haaland G.I., Peel C.J., Bauman D.E. and Steinhour W.D., Administration of bovine growth hormone to high yielding Holstein cows. I. Influence on in vivo energy metabolism. *Journal of Dairy Science.*, 65(1): 120-125 (2008).
5. Stelwagen K., Davis S.R., Farr V.C., Prosser C.G. and Sherlock R.A., Mammary epithelial cell tight junction integrity and mammary blood flow during an extended milking interval in goats. *Journal of Dairy Science.*, 77: 426-432 (2010).
6. Thokal H.E., Kiser T. and Loewenstein M., Influence of milking frequency on productive and reproductive efficiencies of dairy cows. *Journal of Dairy Science.*, 68: 732-742 (2014).
7. Wilde C.J., Addey C.V.P., Boddy L.M. and Peaker M., Autocrine regulation of milk secretion by protein in milk. *Biochemistry Journal.*, 305: 51-58 (2010).
8. Tulloh N.M., Physical studies of the alimentary tract of grazing cattle. IV. Dimensions of the tract in lactating and non-lactating cows. *New Zealand Journal of Agriculture Research.*,

9: 999-1008 (2016).

9. Negrao J.A., Marnet P.G. and Labussiere J., Effect of milking frequency on oxytocin release and milk production in dairy ewes. *Small Ruminant Research.*, 39: 181-187 (2011).

10. Pearson R.E., Fulton L.A., Thompson P.D. and Smith J.W., Three times a day milking during the first half of lactation. *Journal of Dairy Science.*, 62: 1941-1950 (2007).

11. Meyer U., Everinghoff M., Gädeken D. and Flachowsky G., Investigations on the water intake of lactating dairy cows. *Livestock Production Science.*, 90: 117-121 (2004).

12. Meyer J., Benlamlih S. and Dahlborn K., Effect of dehydration, rehydration and hyperhydration in the black Moroccan goat. *Crop Biochemistry and Animal Physiology.*, 109: 1017-1026 (2014).

13. Little W., Sansom B.F., Manston R. and Allen W.M., Effects of restricting the water intake of dairy cows upon their milk yield, body weight and blood composition. *Journal Animal Production.*, 22: 329-339 (2012).

14. Li P., Rudland P.S., Fernig D.G., Finch L.M.B. and Wilde C.J., Modulation of mammary development and programmed cell death by the frequency of milk removal in lactating goats. *Journal of Physiology.*, 519: 885-900 (2009).

15. Linzell J.L., Measurement of udder volume in live goats as an index of mammary growth and function. *Journal of Dairy Science.*, 49: 307-311 (2016).

16. King K.R. and Stockdale C.R., Milk yield of dairy cows given restricted access to water in a Mediterranean-type climate. *Animal Production Science.*, 21: 167-171 (2014).

17. Knight C.H., Hillerton J.E., Kerr M.A., Teverson R.M., Turvey A. and Wilde C.J., Separate and additive stimulation of bovine milk yield by the local and systemic galactopoietic stimuli of frequent milking and growth hormone. *Journal of Dairy Research.*, 59: 243-252 (2012).

18. Hale, S.A., Capuco, A.V. and Erdman, R.A., Milk yield and mammary growth effects due to increased milking frequency during early lactation. *Journal of Dairy Science.*, 86: 2061-2071 (2013).

19. Jurjanz S., Laurent F. and Graupner M., Einfluß einer erhöhten Melkfrequenz auf die Milchzusammensetzung. *Mh. Veterinary Medicine.*, 48: 631-634 (2009).

20. Khan M.A., Nisa M. and Sarwar M., Techniques measuring digestibility for the nutritional evaluation of feeds. *International Journal Agriculture & Biology.*, 5: 91-94 (2003).

21. Khan R., Qureshi M., Mushtaq S., Ghufraulah A. and Naveed A., Effect of quality and frequency of drinking water on productivity and fertility

of dairy buffaloes. *J. Animal Plant Science.*, 22: 96-101 (2012).

22. Hansen L.B., Young C.W., Miller K.P. and Touchberry R.W., Health care requirements of dairy cattle. I. Response to milk yield selection. *Journal Dairy Science.*, 62: 1922-1930 (2007).

23. Hilali N.E., Maltz A., Halevi M. and Shinder D., Metabolism of water, sodium, potassium and chlorine by high yielding dairy cows at the onset of lactation. *Journal Dairy Science.*, 80: 949-956 (2015).

24. Carruthers V.R., Davis S.R., Bryant A.M., Henderson H.V., Morris C.A. and Copeman P.J.A. Response of Jersey and Freisian cows to once a day milking and prediction of response based on udder characteristics and milk composition. *Journal of Dairy Research.*, 60: 1-11 (2013).

25. Chamberlain, A., Milk production in the tropics. Longman Scientific and Technical. England. P. 242 (2011).

26. Bauman D.E., Akers R.M., Chapin L.T., Tucker H.A. and Convey E.M., Effect of level of intake on serum concentrations of prolactin and growth hormone in lactating cows. *Journal of Dairy Science.*, 62(1): 114-116 (2008).

27. Archer P., Milking three times a day. Milk Marketing Board Report, 34. Farm Management Service (FMI). Great Britain. p. 33 (2013).

28. Aspmisra A.K. and Singh K., Effect of water deprivation on dry matter intake, nutrient utilization and metabolic water production in goats under semi-arid zone of India. *Small Ruminant Research.*, 46: 159-165 (2002).

29. Banerjee, G.C., A textbook of animal husbandry. 8th Edi. Oxford and IBH publishing Co. Pvt. Ltd. New Dehli (2009).

30. Adogla-Bessa T. and Aganga A.A., Responses of Tswana goats to various lengths of water deprivation. *South African Journal of Animal Science.*, 30: 87-91 (2000).

31. Amos SR., Farr, V.C. and Stelwagen K., Regulation of yield loss and milk composition during once-daily milking: a review. *Livestock Production Science.*, 59: 77-94 (2015).

32. Andrew T., Alamer M. and Al-Hozab A., Effect of water deprivation and season on feed intake, body weight and thermoregulation in Awassi and Najdi sheep breeds in Saudi Arabia. *Journal of Arid Environment.*, 59: 71-84 (2014).

33. Barnes M.A., Pearson R.E. and Luke-Wilson A.J., Effect of milking frequency and selection for milk yield on productive efficiency of Holstein cows. *Journal of Dairy Science.*, 73: 1603-1611 (2010).

34. Blake R.W. and Custodio A.A., Feed efficiency: a composite trait of dairy cattle.

Journal Dairy Science., 67: 2075-2080 (2014).

35. Dahlborn K., Effect of milking frequency on mammary functioning and shape of the lactation curve. Journal of Dairy Science., 84: 204-211 (2011).
36. DePeters E.J., Smith N.E. and Acedo-Rico J., Three or two times daily milking of older cows and first lactation cows for entire lactation. Journal of Dairy Science., 68: 123-132 (2015).
37. Hadjigeorgiou I., Dardamani K., Goulas C. and Zervas G., The effect of water availability on feed intake and digestion in sheep. Small Ruminant Research., 37: 147-150 (2000).
38. Devendra C., The comparative efficiency of feed utilization of ruminants in the tropics. Tropical Science., 13: 123-132 (2014).
39. Capote J., Lopez J.L., Caja G., Peris S., Arguello A. and Darmanin N., The effects of milking once or 2x daily throughout lactation on milk production of Canarian dairy goats. In: Milking and milk production of dairy sheep and goats. Barillet, F. and Zervas, N.P. (ed.). Wageningen, Netherlands. p. 267-273 (2009).
40. Alamer M., Effect of water restriction on lactation performance of Aardi goats under heat stress conditions. Small Ruminant Research., 84: 76-81(2015).
41. Alamer M. and Al-Hozab A., Effect of water deprivation and season on feed intake, body weight and thermoregulation in Awassi and Najdi sheep breeds in Saudi Arabia. Journal of Arid Environment., 59: 71-84 (2004).
42. Bell A.W., Regulation of organic nutrient metabolism during transition from late pregnancy to early lactation. Journal of Animal Science., 73: 2804-2819 (2011).
43. Erdman R.A. and Varner M., Fixed yield responses to increased milking frequency. Journal Dairy Science., 78: 1199-1203 (2015).
44. Mengistu U., Dahlborn K. and Olsson K., Effects of intermittent watering on water balance and feed intake in male Ethiopian Somali goats. Small Ruminant Research., 67: 45-54 (2007).
45. Poole D.A., The effects of milking cows. Archivos de zootecnia., 61: 465-470 (2011).
46. Senn M.R., Patil V.C. and Udar S.A., Effect of drinking water frequency on milk yield, fat, total solids and solids-not-fat content in crossbred cows. Indian Journal of Animal Research., 38: 47-49 (2016).
47. Senn M., Gross-Luem S.I.N.A., Kaufmann A. and Langhans W., Effect of water deprivation on eating patterns of lactating cows fed grass and corn pellets ad libitum. Physiological Behavior, 60: 1413-1418 (2016).
48. Peel C.J., Fronk T.J., Bauman D.E. and Gorewit R.C., Effect of exogenous growth hormone in early and late lactation on lactational performance

of dairy cows. Journal of Dairy Science., 66: 776-786 (2013).

49. Aganga U., Dahlborn K. and Olsson K., Effects of intermittent watering on water balance and feed intake in male Ethiopian Somali goats. Small Ruminant Research., 67: 45-54 (2012).
50. Burgos M.S., Senn M., Sutter F., Kreuzer M. and Langhans W., Effect of water restriction on feeding and metabolism in dairy cows. American Journal of Physiology., 280: 418-427 (2001).
51. Burgos M.S., Senn M., Sutter F., Kreuzer M. and Langhans W., Effect of water restriction on feeding and metabolism in dairy cows. American Journal of Physiology., 28(1): 418-427 (2015).
52. Royle C., Gamsworth P.C., McArthur A.J. and Mephram T.B., Effect of frequent milking on heart rate and other physiological variables in dairy cows. In: Int. Symp. 46. Prospects for Automatic Milking, Hrdoc. Wageningen. The Netherlands. p. 237 (2015).
53. Salama A.A.K, Such X., Caja G., Rovai M., Casals R., Albanell E., Marn M.P. and Mart A., Effects of once versus 2x daily milking throughout lactation on milk yield and milk composition in dairy goats. Journal of Dairy Science., 86: 1673-1680 (2013).
54. Stockdale E., Stelwagen K. and Knight C.H., Effect of unilateral once or 2x daily milking of cows on milk yield and udder characteristics in early and late lactation. Journal of Dairy Research., 64: 487-494 (2011).